

Breakdown of a Magnetization Plateau due to Anisotropy in Heisenberg Mixed-Spin Chains

Shoji Yamamoto

Department of Physics, Okayama University, Tsushima, Okayama 700-8530, Japan

Tôru Sakai

Faculty of Science, Himeji Institute of Technology, Ako, Hyogo 678-1297, Japan

(18 March 1999)

We discuss the critical behavior of the spin- $(1, \frac{1}{2})$ Heisenberg ferrimagnetic chain in a magnetic field, whose magnetization curve exhibits a plateau at a third of the full magnetization. A bond alternation stabilizes the massive state, whereas an exchange anisotropy causes the breakdown of the plateau and the onset of a gapless spin-fluid state, where the transition, lying in the XY but ferromagnetic region, is of Kosterlitz-Thouless type. In order to elucidate significant quantum effects, we investigate the model of classical version as well.

PACS numbers: 75.10.Jm, 75.40.Mg, 75.30.Kz

I. INTRODUCTION

Ground-state magnetization curves of quantum spin chains have been attracting much current interest due to their quantized plateaux as functions of a magnetic field. Several years ago Hida [1] revealed that a spin- $\frac{1}{2}$ ferromagnetic-antiferromagnetic-antiferromagnetic trimerized chain exhibits a plateau in its magnetization curve at a third of the full magnetization. Although it was already familiar that, in the presence of a field, integer-spin Heisenberg antiferromagnetic chains remain massive from zero field up to a critical field [2], yet the magnetization plateau at a fractional value of the full magnetization was still met with a surprise. Since then various low-dimensional quantum spin systems in a field have been investigated, including polymerized spin chains [3–8], spin chains with anisotropy [9] or four-spin exchange coupling [10], and decorated spin ladders [11–13]. Experimental observations [14,15] of quantized magnetization plateaux have also been reported. In such circumstances, generalizing the Lieb-Schultz-Mattis theorem [16,17], Oshikawa, Yamanaka, and Affleck (OYA) [18] found a criterion for the *fractional quantization*. They pointed out that quantized plateaux in magnetization curves may appear under the condition

$$S_{\text{unit}} - m = \text{integer}, \quad (1.1)$$

where S_{unit} is the sum of spins over all sites in the unit period and m is the magnetization M divided by the number of the unit cells.

Mixed-spin chains are the system of all others that stimulates us in this context. There exists a large amount of chemical knowledge [19] on quantum ferrimagnets. In an attempt to realize a quasi-one-dimensional ferrimagnetic system, Gleizes and Verdager [20] synthesized a few bimetallic compounds such as $\text{AMn}(\text{S}_2\text{C}_2\text{O}_2)_2(\text{H}_2\text{O})_3 \cdot 4.5\text{H}_2\text{O}$ ($A = \text{Cu, Ni, Pd, Pt}$).

Then numerous chemical explorations [21,22] followed and various examples of a ferrimagnetic one-dimensional compound were systematically obtained. The vigorous experimental research motivated theoretical investigations into Heisenberg ferrimagnets. Drillon *et al.* [23] pioneeringly carried out numerical diagonalizations of spin- $(S, \frac{1}{2})$ Heisenberg Hamiltonians for $S = 1$ to $\frac{5}{2}$ and revealed typical thermodynamic properties of ferrimagnetic mixed-spin chains. In recent years, quantum ferrimagnets have met with further theoretical understanding [24–34] owing to various tools such as field [24,34] and spin-wave [25,26,32,33] theories, matrix-product formalism [27,28], and quantum Monte Carlo [26,29,32] and density-matrix renormalization-group [25,30,32] techniques. In particular, their mixed nature, showing both ferromagnetic and antiferromagnetic aspects [32], has lately attracted considerable attention.

However, little is known about quantum ferrimagnetic behavior in a magnetic field [27], especially about magnetization curves [31]. Although anisotropy is an interesting and important factor from an experimental point of view, there exist few arguments on anisotropic models in a field. Now, considering the OYA argument and the accumulated chemical knowledge on ferrimagnetic compounds, the magnetization process of realistic mixed-spin-chain models arouses our interest all the more and indeed deserves urgent communication. In an attempt to serve as guides for further experimental study, we here consider an alignment of alternating spins S and s in a field, as described by the Hamiltonian

$$\mathcal{H} = \sum_{j=1}^N [(\mathbf{S}_j \cdot \mathbf{s}_j)_\alpha + \delta(\mathbf{s}_j \cdot \mathbf{S}_{j+1})_\alpha - H(S_j^z + s_j^z)], \quad (1.2)$$

where $(\mathbf{S} \cdot \mathbf{s})_\alpha = S^x s^x + S^y s^y + \alpha S^z s^z$. We note that even the bond alternation δ is now experimentally adjustable [22]. According to the OYA criterion (1.1), as H increases from zero to the saturation field

$$H_{\text{sat}} = \frac{1}{2}(1 + \delta) \left[\alpha(S + s) + \sqrt{\alpha^2(S - s)^2 + 4Ss} \right], \quad (1.3)$$

the model (1.2) may exhibit quantized plateaux at $m = \frac{1}{2}$ (1), $\frac{3}{2}$ (2), $\dots$, $S + s - 1$. Though a multi-plateau problem is a fascinating subject, we restrict our argument to the simplest case of $(S, s) = (1, \frac{1}{2})$ in the following. This is, on the one hand, because we at first aim at understanding the typical and essential behavior of quantum ferrimagnets in a field, and, on the other hand, because the low-energy structure of the model (1.2) remains qualitatively the same [24,32] as long as $S \neq s$. Then, a plateau is expected at $m = \frac{1}{2}$. At the Heisenberg point, the ground state of the Hamiltonian (1.2) without field is a multiplet of spin $N/2$ [35]. The ferromagnetic excitations, reducing the ground-state magnetization, exhibit a gapless dispersion relation, whereas the antiferromagnetic ones, enhancing the ground-state magnetization, are gapped from the ground state [29]. Therefore, at the isotropic point, m jumps up to $\frac{1}{2}$ just as a field is applied and forms a plateau for $H_{c1} \leq H \leq H_{c2}$ [31], where H_{c1} and H_{c2} are the lower and upper critical fields, equal to 0 and the antiferromagnetic gap, respectively.




FIG. 1. Schematic view of the low-energy structure of the spin-(1, $\frac{1}{2}$) quantum ferrimagnetic chain with anisotropic exchange coupling near the Heisenberg point $\alpha = 1$: (a) the Ising region $\alpha > 1$ and (b) the XY region $\alpha < 1$.

In the presence of exchange anisotropy, the above argument should be modified, where the $(N + 1)$ -fold degenerate ground-state multiplet splits [24,30], as illustrated in Fig. 1. In the Ising region, the ground state is a doublet of $M = \pm N/2$ and therefore H_{c1} remains 0. As α increases, H_{c2} comes to be given as $(1 + \delta)\alpha$ and the magnetization curve ends up with a trivial step. Thus we take little interest in this region. In the XY region, on the other hand, the ground state is a singlet of $M = 0$. Now H_{c1} moves away from 0 and the plateau shrinks as α decreases (see Fig. 2 below). Here arises a stimulative problem: how stable the plateau is against the anisotropy and what comes over the plateau phase? In this article, we demonstrate that the plateau survives the XY anisotropy in the entire antiferromagnetic region

and vanishes in the ferromagnetic region. The transition is of Kosterlitz-Thouless (KT) type [36] and a gapless spin-fluid phase [37] appears instead.




FIG. 2. The ground-state magnetization curves for the quantum Hamiltonian (1.2) at various values of α : (a) $\delta = 1$ and (b) $\delta = 0.6$.




FIG. 3. (a) Scaled quantity $N\Delta_N$ versus α at $\delta = 1$ and $\delta = 0.6$. (b) The central charge c and the critical exponent η versus α in the vicinity of the phase boundary at $\delta = 1$ and $\delta = 0.6$.

phase boundary is also shown in Fig. 4 by a dotted line, which is somewhat discrepant from the highly accurate estimate based on η . The PRG equation applied to gapful-to-gapful phase transitions yields an accurate solution, to be sure, but, for transitions to a gapless phase, including those of KT type, the PRG analysis is likely to miss the correct solution due to essential corrections to the scaling law (2.3), overestimating the gapful-phase region [44,45]. The present PRG solution may still be recognized as the lower boundary of α_c .




FIG. 5. The ground-state magnetization curves for the classical Hamiltonian (1.2) with $\delta = 1$ at various values of α .

III. SUBLATTICE MAGNETIZATIONS

In an attempt to elucidate how much effect quantum fluctuations have on the stability of the plateau, we investigate the Hamiltonian (1.2) of classical version as well, where $\mathbf{S}_j$ and $\mathbf{s}_j$ are classical vectors of magnitude 1 and $\frac{1}{2}$, respectively. We show in Fig. 5 the classical magnetization curves. We note that the classical model also exhibits a plateau at $m = \frac{1}{2}$. The magnetization curves in the Ising region are not so far from the quantum behavior, though we have not shown them explicitly. However, the classical plateau can hardly stand the anisotropy of XY type. In this context, it is interesting to observe sublattice magnetizations separately. We show in Fig. 6 the configuration of each classical spin as a function of a field. The classical plateau is nothing but a Néel-ordered state. In other words, without the fully ordered staggered magnetization, classical spins could not form a magnetization plateau. On the other hand, Fig. 7 shows that quantum spins can form a magnetization plateau with any combination of sublattice magnetizations. It is the case with the quantum model as well that sublattice magnetizations themselves freeze while going through the plateau. However, as long as the XY exchange interaction exists, they are in general reduced from the full values 1 and $-\frac{1}{2}$,

respectively. It is quantum fluctuations that stabilize the plateau with unsaturated sublattice magnetizations.




FIG. 6. The ground-state sublattice magnetizations per unit cell as functions of a field for the classical Hamiltonian (1.2) with $\delta = 1$ at various values of α : (a) the larger spin $S = 1$ and (b) the smaller spin $\frac{1}{2}$.




FIG. 7. The ground-state sublattice magnetizations per unit cell as functions of a field for the quantum Hamiltonian (1.2) with $\delta = 1$ at various values of α : (a) the larger spin $S = 1$ and (b) the smaller spin $\frac{1}{2}$.

One more interesting observation on the quantum spin configuration is that the collapse of the staggered order

where N_{occ} is the number of occupied states, we are allowed to expect magnetization plateaux at $m = \pm \frac{1}{2}$. The inclusion of the bond alternation δ results in the enhancement of the gap, which is consistent with Fig. 2. Qualitatively the same scenario is available for a ferromagnetic-ferromagnetic-antiferromagnetic trimerized chain, as was pointed out by two pioneering authors [48,49]. The present analysis is not strictly comparable to the original argument unless $\alpha = 1$. However, the nonvanishing gap in the $\gamma \rightarrow \infty$ limit may be a qualitative evidence for the existence of the plateau at the XY point in the original model (1.2). We further show in Fig. 9 the sublattice magnetizations in the ground state of the replica model with $\alpha = 0$ as functions of a field at a few values of $\gamma > 1$. We are convinced all the more that the Néel order has already disappeared and both the spins 1 and $\frac{1}{2}$ have the same-sign z components at the XY point.

In recent years, a massive-to-spin-fluid phase transition of KT type has been given a great deal of attention [50–56] in the context of Haldane’s conjecture [2]. In such cases the critical point never goes beyond the XY point. The magnetization plateau in our argument should be distinguished from the gap immediately above the ground state, to be sure, but, compared with Haldane’s scenario [42], the present observation looks novel and is fascinating to be further studied. There may be a new mass-generation mechanism peculiar to quantum mixed-spin chains, other than the valence-bond picture [57]. Quite recently Okamoto and Kitazawa [58] have reported that the magnetization plateau of the spin- $\frac{1}{2}$ trimerized chain which is closely related with the present model also disappears in the XY ferromagnetic region. We hope that our investigation, combined with such an argument from a different viewpoint, will contribute toward revealing the possibly novel scenario for the breakdown of quantized plateaux.

ACKNOWLEDGMENTS

It is a pleasure to thank H.-J. Mikeska and U. Schollwöck for helpful discussions. This work was supported by the Japanese Ministry of Education, Science, and Culture through Grant-in-Aid No. 09740286 and by the Okayama Foundation for Science and Technology. The numerical computation was done in part using the facility of the Supercomputer Center, Institute for Solid State Physics, University of Tokyo.

-
- [1] K. Hida, J. Phys. Soc. Jpn. **63**, 2359 (1994).
 - [2] F. D. M. Haldane, Phys. Lett. **93A**, 464 (1983); Phys. Rev. Lett. **50**, 1153 (1983).
 - [3] K. Okamoto, Solid State Commun. **98**, 245 (1995).
 - [4] T. Tonegawa, T. Nakao, and M. Kaburagi, J. Phys. Soc. Jpn. **65**, 3317 (1996).
 - [5] K. Totsuka, Phys. Lett. A **228**, 103 (1997); Phys. Rev. B **57**, 3454 (1998).
 - [6] H. Nakano and M. Takahashi, J. Phys. Soc. Jpn. **67**, 1126 (1998).
 - [7] D. C. Cabra and M. D. Grynberg, Phys. Rev. B **59**, 119 (1999);
 - [8] A. Honecker, Phys. Rev. B **59**, 6790 (1999).
 - [9] T. Sakai and M. Takahashi, Phys. Rev. B **57**, 3201 (1998).
 - [10] T. Sakai and Y. Hasegawa, preprint (cond-mat/9809291).
 - [11] D. C. Cabra, A. Honecker, and P. Pujol, Phys. Rev. Lett. **79**, 5126 (1997); Phys. Rev. B **58**, 6241 (1998).
 - [12] K. Tandon, S. Lal, S. K. Pati, S. Ramasesha, and D. Sen, Phys. Rev. B **59**, 396 (1999).
 - [13] A. K. Kolezhuk, Phys. Rev. B **59**, February 1 (1999).
 - [14] Y. Narumi, M. Hagiwara, R. Sato, K. Kindo, H. Nakano, and M. Takahashi, Physica B **246-247**, 509 (1998).
 - [15] W. Shiramura, K. Takatsu, B. Kurniawan, H. Tanaka, H. Uekusa, Y. Ohashi, K. Takizawa, H. Mitamura, and T. Goto, J. Phys. Soc. Jpn. **67**, 1548 (1998).
 - [16] E. Lieb, T. Schultz, D. Mattis, Ann. Phys. **16**, 407 (1961).
 - [17] I. Affleck, Phys. Rev. **37**, 5186 (1988).
 - [18] M. Oshikawa, M. Yamanaka, and I. Affleck, Phys. Rev. Lett. **78**, 1984 (1997).
 - [19] O. Kahn, *Magnetism of the heteropolymetallic systems*, Structure and Bonding **68**, 91 (Springer-Verlag, 1987); O. Kahn, Y. Pei, and Y. Journaux, in *Inorganic Materials*, edited by D. W. Bruce and D. O’Hare (Wiley, New York, 1995), p. 95.
 - [20] A. Gleizes and M. Verdaguer, J. Am. Chem. Soc. **103**, 7373 (1981); *ibid.* **106**, 3727 (1984).
 - [21] Y. Pei, M. Verdaguer, O. Kahn, J. Sletten, and J.-P. Renard, Inorg. Chem. **26**, 138 (1987); O. Kahn, Y. Pei, M. Verdaguer, J.-P. Renard, and J. Sletten, J. Am. Chem. Soc. **110**, 782 (1988); P. J. van Koningsbruggen, O. Kahn, K. Nakatani, Y. Pei, J.-P. Renard, M. Drillon, and P. Legoll, Inorg. Chem. **29**, 3325 (1990).
 - [22] Y. Pei, O. Kahn, J. Sletten, J.-P. Renard, R. Georges, J.-C. Gianduzzo, J. Curely, and Q. Xu, Inorg. Chem. **27**, 47 (1988).
 - [23] M. Drillon, J. C. Gianduzzo, and R. Georges, Phys. Lett. **96A**, 413 (1983); M. Drillon, E. Coronado, R. Georges, J. C. Gianduzzo, and J. Curely, Phys. Rev. B **40**, 10992 (1989).
 - [24] F. C. Alcaraz and A. L. Malvezzi, J. Phys. A **30**, 767 (1997).
 - [25] S. K. Pati, S. Ramasesha, and D. Sen, Phys. Rev. B **55**, 8894 (1997); J. Phys.: Condens. Matter **9**, 8707 (1997).
 - [26] S. Brehmer, H.-J. Mikeska, and S. Yamamoto, J. Phys.: Condens. Matter **9**, 3921 (1997).
 - [27] A. K. Kolezhuk, H.-J. Mikeska, and S. Yamamoto, Phys. Rev. B **55**, 3336 (1997); K. Maisinger, U. Schollwöck, S. Brehmer, H.-J. Mikeska, and S. Yamamoto, Phys. Rev. B **58**, 5908 (1998); A. K. Kolezhuk, H.-J. Mikeska, K. Maisinger, and U. Schollwöck, preprint (cond-mat/9812326).

- [28] H. Niggemann, G. Uimin, and J. Zittartz, J. Phys.: Condens. Matter **9**, 9031 (1997); *ibid.* **10**, 5217 (1998).
- [29] S. Yamamoto, Int. J. Mod. Phys. C **8**, 609 (1997); S. Yamamoto, S. Brehmer, and H.-J. Mikeska, Phys. Rev. B **57**, 13610 (1998); S. Yamamoto and T. Sakai, J. Phys. Soc. Jpn. **67**, 3711 (1998).
- [30] T. Ono, T. Nishimura, M. Katsumura, T. Morita, and M. Sugimoto, J. Phys. Soc. Jpn. **66**, 2576 (1997).
- [31] T. Kuramoto, J. Phys. Soc. Jpn. **67**, 1762 (1998).
- [32] S. Yamamoto and T. Fukui, Phys. Rev. B **57**, 14008 (1998); S. Yamamoto, T. Fukui, K. Maisinger, and U. Schollwöck, J. Phys.: Condens. Matter **10**, 11033 (1998); S. Yamamoto, Phys. Rev. B **59**, 1024 (1999).
- [33] N. B. Ivanov, Phys. Rev. B **57**, 14024 (1998).
- [34] M. Abolfath, H. Hamidian, and A. Langari, preprint (cond-mat/9901063).
- [35] E. Lieb and D. Mattis, J. Math. Phys. **3**, 749 (1962).
- [36] J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973).
- [37] F. D. M. Haldane, Phys. Rev. B **25**, 4925 (1982).
- [38] T. Sakai and M. Takahashi, Phys. Rev. B **43**, 13383 (1991);
- [39] D. Shanks, J. Math. Phys. **34**, 1 (1955).
- [40] J. L. Cardy, J. Phys. A **17**, L385 (1984); H. W. Blöte, J. L. Cardy and M. P. Nightingale, Phys. Rev. Lett. **56**, 742 (1986).
- [41] I. Affleck, Phys. Rev. Lett. **56**, 746 (1986).
- [42] H. J. Schulz, Phys. Rev. B **34**, 6372 (1986).
- [43] M. P. Nightingale, Physica **83A**, 561 (1976).
- [44] K. Nomura and K. Okamoto, J. Phys. Soc. Jpn. **62**, 1123 (1993).
- [45] K. Okamoto and K. Nomura, J. Phys. A: Math. Gen. **29**, 2279 (1996).
- [46] S. Yamamoto and T. Sakai, unpublished.
- [47] See for example, K. Hida, Phys. Rev. B **45**, 2207 (1992); *ibid.* **46**, 8268 (1992).
- [48] C. E. Zaspel, J. Chem. Phys. **86**, 4713 (1987).
- [49] K. Okamoto, Solid State Commun. **83**, 1039 (1992).
- [50] R. Botet, R. Jullien, and M. Kolb, Phys. Rev. B **28**, 3914 (1983).
- [51] J. Solyom and T. Ziman, Phys. Rev. B **30**, 3980; H. J. Schulz and T. Ziman, *ibid.* **33**, 6545 (1986).
- [52] F. C. Alcaraz and A. Moreo, Phys. Rev. B **46**, 2896 (1992).
- [53] U. Schollwöck and Th. Jolicœur, Europhys. Lett. **30**, 493 (1995).
- [54] M. Yajima and M. Takahashi, J. Phys. Soc. Jpn. **63**, 3634 (1994).
- [55] A. L. Malvezzi and F. C. Alcaraz, J. Phys. Soc. Jpn. **64**, 4485 (1995).
- [56] J. P. Neirotti and M. J. de Oliveira, Phys. Rev. B **59**, 3303 (1999).
- [57] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987); Commun. Math. Phys. **115**, 477 (1988).
- [58] K. Okamoto and A. Kitazawa, preprint (cond-mat/9809138).

TABLE I. The antiferromagnetic excitation gap Δ as a function of δ at the Heisenberg point $\alpha = 1$.

δ	1	0.8	0.6	0.4	0.2	0
Δ	1.7591	1.6042	1.4986	1.4500	1.4558	$\frac{3}{2}$